| Туре | | | | Resistance ✓✓ = Good ✓ = Fair x = Not | | | | |
|-----------|---------------------------|------|-------|--|------------|--|--|--|
| 1st Early | 2nd Early | Main | Salad | Blight | Scab | Eelworm | Balga Baldock Allotment and Leisure Gardeners Association | |
| • | | | | ✓ | × | √ | Very early with high dry matter. Can be dug up early and eaten as a salad potato or left to mature to baking size potatoes. | |
| • | | | | ✓ | * * | × | One of the most popular varieties. Very variable in size and shape. Good yield and flavour but needs reasonable soil condition. Good for boiling | 90 |
| • | | | | 1 | * * | ✓ | White skin with creamy flesh produces high number of tubers. Good for boiling and steaming | 20 |
| • | | | | * | * * | √ | Excellent all-rounder with light yellow skin and flesh. Has a good resistance to tuber blight. ORGANIC so ideal for the organic grower. | 20 |
| • | | | | ✓ | * | × | Very good cooking qualities and a gardener's favourite for flavour. Good for boiling | 90 |
| • | | | | ~ | * * | 44 | Suitable for early French fries, with medium yields, early bulking. Susceptible to late blight on foliage, Blackleg and potato virus Yo. Resistance to bruising and splitting. Susceptibility to potato cyst nematode. | 3 |
| • | | | | * | ~ | 11 | Possibly the earliest early with a very mild flavour. Produces huge crops of uniform attractive white tubers. Has double eelworm resistance, should not be left in ground. Good for boiling | 80 |
| • | | | | ~ | * * | ✓ | Renowned for their boiling qualities when cooked straight after harvest. Excellent drought resistance, low resistance to late blight on foliage and late blight on tubers, dry rot, potato leafroll virus and potato virus Yo. | 20 |
| | 1st Early • • • • • • • • | | | | Type Salad | Type Scab Salad Main 1st Early 1st Early Main Ma | Type | Type Fair Fa |

Table content based on information in the Kings Seeds catalogue and the AHDB website